

7.2: CELL CYCLE AND CELL DIVISION

□ SO MANY CELLS!

The baby in Figure 7.2.1 has a lot of growing to do before they are as big as their mom. Most of their growth will be the result of cell division. By the time the baby is an adult, their body will consist of trillions of cells. Cell division is just one of the stages that all cells go through during their life. This includes cells that are harmful, such as cancer cells. Cancer cells divide more often than normal cells and grow out of control. In fact, this is how cancer cells cause illness. In this concept, you will read about how cells divide, what other stages cells go through, and what causes cancer cells to divide out of control and harm the body.



Figure 7.2.1: Baby and mother

THE CELL CYCLE

Cell division is the process in which one cell, called the parent cell, divides to form two new cells, referred to as daughter cells. How this happens depends on whether the cell is prokaryotic or eukaryotic. Cell division is simpler in prokaryotes than eukaryotes because prokaryotic cells themselves are simpler. Prokaryotic cells have a single circular chromosome, no nucleus, and few other organelles. Eukaryotic cells, in contrast, have multiple chromosomes contained within a nucleus and many other organelles. All of these cell parts must be duplicated and then separated when the cell divides. Cell division is just one of several stages that a cell goes through during its lifetime. The **cell cycle** is a repeating series of events that include growth, DNA synthesis, and cell division. The cell cycle in prokaryotes is quite simple: the cell grows, its DNA replicates, and the cell divides. This form of division in prokaryotes is called asexual reproduction. In eukaryotes, the cell cycle is more complicated.

EUKARYOTIC CELL CYCLE

Figure 7.2.2 represents the cell cycle of a eukaryotic cell. As you can see, the eukaryotic cell cycle has several phases. The mitotic phase (M) includes both mitosis and cytokinesis. This is when the nucleus and then the cytoplasm divide. The other three phases (G_1 , S, and G_2) are generally grouped together as **interphase**. During interphase, the cell grows, performs routine life processes, and prepares to divide. These phases are discussed below.

INTERPHASE

The Interphase of the eukaryotic cell cycle can be subdivided into the following phases (Figure 7.2.2).

- **Growth Phase 1 (G_1):** The cell spends most of its life in the first gap (sometimes referred to as growth) phase, G_1 . During this phase, a cell undergoes rapid growth and performs its routine functions. During this phase, the biosynthetic and metabolic activities of the cell occur at a high rate. The synthesis of amino acids and hundreds of thousands or millions of proteins that are required by the cell occurs during this phase. Proteins produced include those needed for DNA replication. If a cell is not dividing, the cell enters the G_0 phase from this phase.
- **G_0 phase:** The G_0 phase is a resting phase where the cell has left the cycle and has stopped dividing. Non-dividing cells in multicellular eukaryotic organisms enter G_0 from G_1 . These cells may remain in G_0 for long periods of time, even indefinitely, such as with neurons. Cells that are completely differentiated may also enter G_0 . Some cells stop dividing when issues of sustainability or viability of their daughter cells arise, such as with DNA damage or degradation, a process called **cellular senescence**. Cellular senescence occurs when normal diploid cells lose the ability to divide, normally after about 50 cell divisions.
- **Synthesis Phase (S):** Dividing cells enter the Synthesis (S) phase from G_1 . For two genetically identical daughter cells to be formed, the cell's DNA must be copied through DNA replication. When the DNA is replicated, both strands of the double helix are used as templates to produce two new complementary strands. These new strands then hydrogen bond to the template strands and two double helices form. During this phase, the amount of DNA in the cell has effectively doubled, though the cell remains in a diploid state.
- **Growth Phase 2 (G_2):** The second gap (growth) (G_2) phase is a shortened growth period in which many organelles are reproduced or manufactured. Parts necessary for mitosis and cell division are made during G_2 , including **microtubules** used in the mitotic spindle.




Figure 7.2.2: Eukaryotic Cell Cycle. The First Gap (G_1), Synthesis (S), and Second Gap (G_2) phases make up interphase (I). The mitotic phase (yellow M) includes mitosis (purple M) and cytokinesis. During cytokinesis two cells result. Some cells do not divide and they enter into G_0 phase.

MITOTIC PHASE

Before a eukaryotic cell divides, all the DNA in the cell's multiple chromosomes is replicated. Its organelles are also duplicated. This happens in the interphase. Then, when the cell divides (mitotic phase), it occurs in two major steps, called **mitosis** and **cytokinesis**, both of which

are described in greater detail in the concept [Mitotic Phase: Mitosis and Cytokinesis](#).

- The first step in the mitotic phase of a eukaryotic cell is mitosis, a multi-phase process in which the nucleus of the cell divides. During mitosis, the nuclear envelope (membrane) breaks down and later reforms. The chromosomes are also sorted and separated to ensure that each daughter cell receives a complete set of chromosomes.
- The second major step is cytokinesis. This step, which occurs in prokaryotic cells as well, is when the cytoplasm divides and two daughter cells form.

Table 7.2.2: Cell Cycle Summary

State	Name	Description
Quiescent Senescent	Resting phase (G ₀)	A resting phase where the cell has left the cycle and has stopped dividing.
Interphase	1st growth phase (G ₁)	Cells increase in size in G ₁ . Cells perform their normal activities.
	Synthesis phase (S)	DNA replication occurs during this phase.
	2nd growth phase (G ₂)	The cell will continue to grow and many organelles will divide during their phase.
Cell division	Mitosis (M)	Cell growth stops at this stage. Mitosis divides the nucleus into two nuclei, followed by cytokinesis which divides the cytoplasm. Two genetically identical daughter cells result.

CONTROL OF THE CELL CYCLE

If the cell cycle occurred without regulation, cells might go from one phase to the next before they were ready. What controls the cell cycle? How does the cell know when to grow, synthesize DNA, and divide? The cell cycle is controlled mainly by regulatory proteins. These proteins control the cycle by signaling the cell to either start or delay the next phase of the cycle. They ensure that the cell completes the previous phase before moving on. Regulatory proteins control the cell cycle at key checkpoints, which are shown in Figure 7.2.3. There are a number of main checkpoints:

1. The G₁ checkpoint: just before entry into the S phase, makes the key decision of whether the cell is big enough to divide. If the cell is not big enough, it goes into the resting period (G₀)
2. DNA synthesis Checkpoint: The S checkpoint determines if the DNA has been replicated properly.
3. The mitosis checkpoint: This checkpoint ensures that all the chromosomes are properly aligned before the cell is allowed to divide.

The Cell Cycle and the Checkpoints




Figure 7.2.3: Checkpoints in the eukaryotic cell cycle ensure that the cell is ready to proceed before it moves on to the next phase of the cycle.

CANCER AND THE CELL CYCLE

Cancer is a disease that occurs when the cell cycle is no longer regulated. This happens because a cell's DNA becomes damaged. This results in mutations in the genes that regulate the cell cycle. Damage can occur due to exposure to hazards such as radiation or toxic chemicals. Cancerous cells generally divide much faster than normal cells. They may form a mass of abnormal cells called a **tumor** (see Figure 7.2.4). The rapidly dividing cells take up nutrients and space that normal cells need. This can damage tissues and organs and eventually lead to death. When uncontrolled cell division happens in the bone marrow, abnormal and nonfunctional blood cells are produced because the division is happening before the cell is ready for division. In these types of cancer, there is not any evident tumor.




Figure 7.2.4: Gross appearance of an opened large intestine specimen containing an invasive **colorectal carcinoma** and two **adenomatous polyps**. The cancer cells have grown out of control and forming a tumor.

7.3: MITOTIC PHASE - MITOSIS AND CYTOKINESIS

□ DIVIDE AND SPLIT

Can you guess what this colorful image represents? It shows a eukaryotic cell during the process of cell division. In particular, the image shows the nucleus of the cell dividing. In eukaryotic cells, the nucleus divides before the cell itself splits in two; and before the nucleus divides, the cell's DNA is replicated, or copied. There must be two copies of the DNA so that each daughter cell will have a complete copy of the genetic material from the parent cell. How is the replicated DNA sorted and separated so that each daughter cell gets a complete set of genetic material? To answer that question, you first need to know more about DNA and the forms it takes.




Figure 7.3.1: Dividing cell stained with fluorescent dyes. You can see chromosomes in blue and spindles in green.

THE FORMS OF DNA

Except when a eukaryotic cell divides, its nuclear DNA exists as a grainy material called **chromatin**. Only when a cell is about to divide and its

DNA has replicated does DNA condense and coil into the familiar X-shaped form of a **chromosome**, like the one shown in Figure 7.3.2. Because DNA has already replicated, each chromosome actually consists of two identical copies. The two copies of a chromosome are called **sister chromatids**. Sister chromatids are joined together at a region called a **centromere**.




Figure 7.3.2: Chromosome. After DNA replicates, it forms X-shaped chromosomes like the one shown here. 1. Chromatid, 2. Centromere, 3. short arm, 4. long arm. Centromere contains proteins called kinetochores (not shown) where spindles attach during mitosis.

The process in which the nucleus of a eukaryotic cell divides is called **mitosis**. During mitosis, the two sister chromatids that make up each chromosome separate from each other and move to opposite poles of the cell. Mitosis occurs in four phases. The phases are called prophase, metaphase, anaphase, and telophase. They are shown in Figure 7.3.3 and described in detail below.




Figure 7.3.3: Mitosis is the phase of the eukaryotic cell cycle that occurs between DNA replication and the formation of two daughter cells. Mitosis has four substages, prophase, metaphase, anaphase, and telophase.

PROPHASE




Figure 7.3.4: Prophase in later stage is called prometaphase. The spindle starts to form during the prophase of mitosis. The spindles start to attach to the Kinetochores of centromeres of sister chromatids during Prometaphase.

The first and longest phase of mitosis is prophase. During prophase, chromatin condenses into chromosomes, and the nuclear envelope (the membrane surrounding the nucleus) breaks down. In animal cells, the centrioles near the nucleus begin to separate and move to opposite poles of the cell. Centrioles are small organelles found only in eukaryotic cells that help ensure the new cells that form after cell division each contain a complete set of chromosomes. As the centrioles move apart, a spindle starts to form between them. The blue spindle, shown in Figure 7.3.4, consists of fibers made of microtubules.

METAPHASE

During **metaphase**, spindle fibers fully attach to the centromere of each pair of sister chromatids. As you can see in Figure 7.3.5, the sister chromatids line up at the equator, or center, of the cell. The spindle fibers ensure that sister chromatids will separate and go to different daughter cells when the cell divides. Some spindles do not attach to the kinetochore

protein of the centromeres. These spindles are called non-kinetochore spindles that help in the elongation of the cell. This is visible in Figure 7.3.5.




Figure 7.3.5: Chromosomes, consisting of sister chromatids, line up at the equator or middle of the cell during metaphase. The blue lines are spindles, and the orange rectangles at the cell poles are centrioles. Some spindles from the opposing centrioles attach with each other, and some spindles attach to the kinetochores of the sister chromosomes from their respective sides. Each chromosome is attached to two spindles.

ANAPHASE

During **anaphase**, sister chromatids separate and the centromeres divide. The sister chromatids are pulled apart by the shortening of the spindle fibers. This is a little like reeling in a fish by shortening the fishing line. One sister chromatid moves to one pole of the cell, and the other sister chromatid moves to the opposite pole (see Figure 7.3.6). At the end of anaphase, each pole of the cell has a complete set of chromosomes




Figure 7.3.6: Anaphase: Sister chromatids break apart and move to the opposite pole with the help of spindles. The newly separated sister chromatids are called chromosomes now.

TELOPHASE

The chromosomes reach the opposite poles and begin to *decondense* (unravel), relaxing once again into a stretched-out chromatin configuration. The mitotic spindles are depolymerized into tubulin monomers that will be used to assemble cytoskeletal components for each daughter cell. Nuclear envelopes form around the chromosomes, and nucleosomes appear within the nuclear area (see Figure 7.3.7).




Figure 7.3.7: Telophase: The chromosomes decondense, spindles start to disappear, two nuclei form in a cell.

CYTOKINESIS

Cytokinesis is the final stage of cell division in eukaryotes as well as prokaryotes. During cytokinesis, the cytoplasm splits in two and the cell divides. The process is different in plant and animal cells, as you can see in Figure 7.3.8. In animal cells, the plasma membrane of the parent cell pinches inward along the cell's equator until two daughter cells form. In the plant cells, a cell plate forms along the equator of the parent cell. Then, a new plasma membrane and cell wall form along each side of the cell plate.




Figure 7.3.8: Cytokinesis is the final stage of eukaryotic cell division. It occurs differently in animal (left) and plant (right) cells. You can see a microfilament ring forming at the center of the elongated animal cell. This creates a depression called cleavage furrow. This invagination ultimately separates the cell cytoplasm into two cells. A cell plate forms at the center of the elongated plant cell. Then a new plasma membrane and cell wall form along each side of the cell plate.

7.4: MUTATIONS AND CANCER

□ WHAT WOULD HAPPEN IF THIS CYCLE PROCEEDS AT WILL?

Your cells may grow and divide without performing their necessary functions, or without fully replicating their DNA, or without copying their organelles. Probably not much good could come of that. So the cell cycle needs to be highly regulated and tightly controlled. And it is.




Figure 7.4.1: Cell Cycle

CONTROL OF THE CELL CYCLE

How does the cell know when to divide? How does the cell know when to replicate its DNA? How does the cell know when to proceed into mitosis or cytokinesis? The answers to these questions have to do with the control of the cell cycle. But how is the cell cycle controlled or regulated? Regulation of the cell cycle involves processes crucial to the survival of a cell. These include the detection and repair of damage to DNA, as well as the prevention of uncontrolled cell division. Uncontrolled cell division can be deadly to an organism; its prevention is critical for survival.

CYCLINS AND KINASES

The cell cycle is controlled by a number of protein-controlled feedback processes. Two types of proteins involved in the control of the cell cycle are **kinases** and **cyclins**. Cyclins activate kinases by binding to them, specifically they activate **cyclin-dependent kinases (CDK)**. Kinases are enzymes that catalyze the transfer of a phosphate group from ATP to another molecule in a cell. They function as a control switch in many cellular functions, turning a function on or off, and regulating other cellular processes. Many times they are involved in activating a cascade of reactions. Cyclins comprise a group of proteins that are rapidly produced at key stages in the cell cycle. Once activated by a cyclin, CDK enzymes activate or inactivate other target molecules through phosphorylation. It is this precise regulation of proteins that triggers advancement through the cell cycle. Leland H. Hartwell, R. Timothy Hunt, and Paul M. Nurse won the 2001 Nobel Prize in Physiology or Medicine for their discovery of these critical proteins.

WHAT MAKES A CELL CANCEROUS?

Cancer is a disease characterized by a population of cells that grow and divide without respect to normal limits. These cancerous cells invade and destroy adjacent tissues, and they may spread throughout the body. The process by which normal cells are transformed into cancer cells is known

as **carcinogenesis**. This process is also known as oncogenesis or tumorigenesis.

Nearly all cancers are caused by mutations in the DNA of abnormal cells. These mutations may be due to the effects of **carcinogens**, cancer-causing agents such as tobacco smoke, radiation, chemicals, or infectious agents. These carcinogens may act as an environmental “trigger,” stimulating the onset of cancer in certain individuals and not others. Do all people who smoke get cancer? No. Can secondhand smoke increase a nonsmoking person's chance of developing lung cancer? Yes. It also increases a nonsmoking person's chance of developing heart disease.

Complex interactions between carcinogens and an individual's genome may explain why only some people develop cancer after exposure to an environmental trigger and others do not. Do all cancers need an environmental trigger to develop? No. Cancer-causing mutations may also result from errors incorporated into the DNA during replication, or they may be inherited. Inherited mutations are present in all cells of the organism.

ONCOGENES AND TUMOR SUPPRESSOR GENES

Some types of cancer occur because of mutations in genes that control the cell cycle. Cancer-causing mutations most often occur in two types of regulatory genes, called proto-oncogenes and tumor-suppressor genes.

- Proto-oncogenes are genes that normally help cells divide. When a proto-oncogene mutates to become an oncogene, it is continuously active, even when it is not supposed to be. This is like a car's accelerator pedal being stuck at full throttle. The car keeps racing at top speed. In the case of a cell, the cell keeps dividing out of control, which can lead to cancer.
- Tumor suppressor genes are genes that normally slow down or stop cell division. When a mutation occurs in a tumor suppressor gene, it can no longer control cell division. This is like a car without brakes. The car can't be slowed or stopped. In the case of a cell, the cell keeps dividing out of control, which can lead to cancer.

Several Mutations to Cause Cancer

Oncogenes may be growth factors, protein kinases, GTPases or transcription factors. **Growth factors** are naturally occurring substances, usually a protein or steroid hormone, capable of stimulating cellular growth, proliferation, and differentiation. They are important for regulating a variety of cellular processes. Usually, they must bind to an extracellular or intracellular receptor to initiate a cellular reaction.

Typically, a series of several mutations that constitutively activate oncogenes and inactivate tumor suppressor genes is required to transform a normal cell into a cancer cell (Figure 7.4.2). Cells have developed a number of control mechanisms to overcome mutations in proto-oncogenes. Therefore, a cell needs multiple mutations to transform into a cancerous cell. A mutation in one proto-oncogene would not cause cancer, as the effects of the mutation would be masked by the normal control of the cell cycle and the actions of tumor suppressor genes. Similarly, a mutation in one tumor suppressor gene would not cause cancer either, due to the presence of many “backup” genes that duplicate its functions. It is only when enough proto-oncogenes have mutated into oncogenes and enough tumor suppressor genes have been deactivated that the cancerous transformation can begin. Signals for cell growth overwhelm the signals for growth regulation, and the cell quickly spirals out of control. Often, because many of these genes regulate the processes

that prevent most damage to the genes themselves, DNA damage accumulates as one ages.




Figure 7.4.2: How Cancer Develops. Mutations in a tumor suppressor gene allow the proliferation of cells. As many times the cells divide, they acquire more mutation. Some mutations may lead to the inactivation of the DNA repair genes. Also, proto-oncogenes may convert into oncogenes due to mutations. Several other mutations inactivate many tumor suppressor genes. Eventually, a series of mutations in tumor-suppressor genes and proto-oncogenes leads to cancer

Usually, oncogenes are dominant alleles, as they contain gain-of-function mutations. The actions of the mutant allele gene product, many times resulting in a constitutively activated protein, are dominant to the gene product produced by the "normal" allele. Meanwhile, mutated tumor

suppressors are generally recessive alleles, as they contain loss-of-function mutations. A proto-oncogene needs only a mutation in one copy of the gene to generate an oncogene; a tumor suppressor gene needs a mutation in both copies of the gene to render both products defective. There are instances when, however, one mutated allele of a tumor suppressor gene can render the other copy non-functional. These instances result in what is known as a dominant negative effect.

5.2 Chromosomes and Mitosis

Lesson Objectives

- Describe chromosomes and their role in mitosis.
- Outline the phases of mitosis.

Vocabulary

- anaphase
- centromere
- chromatid
- chromatin
- chromosome
- gene
- homologous chromosomes
- metaphase
- prophase
- telophase

Introduction

In eukaryotic cells, the nucleus divides before the cell itself divides. The process in which the nucleus divides is called mitosis. Before mitosis occurs, a cell's DNA is replicated. This is necessary so that each daughter cell will have a complete copy of the genetic material from the parent cell. How is the replicated DNA sorted and separated so that each daughter cell gets a complete set of the genetic material? To understand how this happens, you need to know more chromosomes.

Chromosomes

Chromosomes are coiled structures made of DNA and proteins. Chromosomes are the form of the genetic material of a cell during cell division. During other phases of the cell cycle, DNA is not coiled into chromosomes. Instead, it exists as a grainy material called **chromatin**.

The vocabulary of DNA: chromosomes, chromatids, chromatin, transcription, translation, and replication is discussed at <http://www.youtube.com/watch?v=s9HPNwXd9fk> (18:23).


**MEDIA**

Click image to the left or use the URL below.

URL: <https://www.ck12.org/flx/render/embeddedobject/269>

Chromatids and the Centromere

DNA condenses and coils into the familiar X-shaped form of a chromosome, shown in **Figure 5.6**, only after it has replicated. (You can watch DNA coiling into a chromosome at the link below.) Because DNA has already replicated, each chromosome actually consists of two identical copies. The two copies are called sister **chromatids**. They are attached to one another at a region called the **centromere**. A remarkable animation can be viewed at <http://www.hmi.org/biointeractive/dna-packaging> .

**FIGURE 5.6**

Chromosome. After DNA replicates, it forms chromosomes like the one shown here.

Chromosomes and Genes

The DNA of a chromosome is encoded with genetic instructions for making proteins. These instructions are organized into units called **genes**. Most genes contain the instructions for a single protein. There may be hundreds or even thousands of genes on a single chromosome.

Human Chromosomes

Human cells normally have two sets of chromosomes, one set inherited from each parent. There are 23 chromosomes in each set, for a total of 46 chromosomes per cell. Each chromosome in one set is matched by a chromosome of the same type in the other set, so there are actually 23 pairs of chromosomes per cell. Each pair consists of chromosomes of the same size and shape that also contain the same genes. The chromosomes in a pair are known as **homologous chromosomes**.

Mitosis and Cytokinesis

During mitosis, when the nucleus divides, the two chromatids that make up each chromosome separate from each other and move to opposite poles of the cell. This is shown in **Figure 5.7**. You can watch an animation of the process at the following link: http://www.biology.arizona.edu/Cell_bio/tutorials/cell_cycle/MitosisFlash.html .




FIGURE 5.7

Mitosis is the phase of the eukaryotic cell cycle that occurs between DNA replication and the formation of two daughter cells. What happens during mitosis?

Mitosis actually occurs in four phases. The phases are called prophase, metaphase, anaphase, and telophase. They are shown in **Figure 5.8** and described in greater detail in the following sections.

Prophase


The first and longest phase of mitosis is **prophase**. During prophase, chromatin condenses into chromosomes, and the nuclear envelope, or membrane, breaks down. In animal cells, the centrioles near the nucleus begin to separate and move to opposite poles of the cell. As the centrioles move, a spindle starts to form between them. The spindle, shown in **Figure 5.9**, consists of fibers made of microtubules.

Metaphase

During **metaphase**, spindle fibers attach to the centromere of each pair of sister chromatids (see **Figure 5.10**). The sister chromatids line up at the equator, or center, of the cell. The spindle fibers ensure that sister chromatids will separate and go to different daughter cells when the cell divides.

Anaphase

During **anaphase**, sister chromatids separate and the centromeres divide. The sister chromatids are pulled apart by the shortening of the spindle fibers. This is like reeling in a fish by shortening the fishing line. One sister chromatid

**FIGURE 5.8**

Mitosis in the Eukaryotic Cell Cycle. Mitosis is the multi-phase process in which the nucleus of a eukaryotic cell divides.


moves to one pole of the cell, and the other sister chromatid moves to the opposite pole. At the end of anaphase, each pole of the cell has a complete set of chromosomes.

Telophase


During **telophase**, the chromosomes begin to uncoil and form chromatin. This prepares the genetic material for directing the metabolic activities of the new cells. The spindle also breaks down, and new nuclear membranes form.

Cytokinesis

Cytokinesis is the final stage of cell division in eukaryotes as well as prokaryotes. During cytokinesis, the cytoplasm splits in two and the cell divides. Cytokinesis occurs somewhat differently in plant and animal cells, as shown in **Figure 5.11**. In animal cells, the plasma membrane of the parent cell pinches inward along the cell's equator until two daughter cells form. In plant cells, a cell plate forms along the equator of the parent cell. Then, a new plasma membrane and cell wall form along each side of the cell plate.

**FIGURE 5.9**

Spindle. The spindle starts to form during prophase of mitosis. Kinetochore on the spindle attach to the centromeres of sister chromatids.

**FIGURE 5.10**


Chromosomes, consisting of sister chromatids, line up at the equator or middle of the cell during metaphase.

The phases of mitosis are discussed in the video: http://www.youtube.com/watch?v=LLKX_4DHE3I .

**MEDIA**

Click image to the left or use the URL below.

URL: <https://www.ck12.org/flx/render/embeddedobject/271>

**FIGURE 5.11**

Cytokinesis is the final stage of eukaryotic cell division. It occurs differently in animal (left) and plant (right) cells.

Lesson Summary

- Chromosomes are coiled structures made of DNA and proteins. They form after DNA replicates and are the form in which the genetic material goes through cell division. Chromosomes contain genes, which code for proteins.
- Cell division in eukaryotic cells includes mitosis, in which the nucleus divides, and cytokinesis, in which the cytoplasm divides and daughter cells form.
- Mitosis occurs in four phases, called prophase, metaphase, anaphase, and telophase.

Lesson Review Questions

Recall

1. What are chromosomes? When do they form?
2. Identify the chromatids and the centromere of a chromosome.
3. List the phases of mitosis.
4. What happens during prophase of mitosis?
5. During which phase of mitosis do sister chromatids separate?
6. Describe what happens during cytokinesis in animal cells.

Apply Concepts

7. If a cell skipped metaphase during mitosis, how might this affect the two daughter cells?

Think Critically

8. Explain how chromosomes are related to chromatin. Why are chromosomes important for mitosis?
9. Explain the significance of the spindle in mitosis.

Points to Consider

Cell division occurs not only as organisms grow. It also occurs when they reproduce.

- What role do you think cell division plays when prokaryotes such as bacteria reproduce?
- How do you think cell division is involved in the reproduction of eukaryotes such as humans?